

Topical Review

Muscle-derived interleukin-6: possible biological effects

Bente Klarlund Pedersen^{*†}, Adam Steensberg^{*†} and Peter Schjerling^{*}

^{*}*The Copenhagen Muscle Research Centre and* [†]*Department of Infectious Diseases, University of Copenhagen, Denmark*

(Received 15 June 2001; accepted after revision 24 August 2001)

Interleukin-6 (IL-6) is produced locally in working skeletal muscle and can account for the increase in plasma IL-6 during exercise. The production of IL-6 during exercise is related to the intensity and duration of the exercise, and low muscle glycogen content stimulates the production. Muscle-derived IL-6 is released into the circulation during exercise in high amounts and is likely to work in a hormone-like fashion, exerting an effect on the liver and adipose tissue, thereby contributing to the maintenance of glucose homeostasis during exercise and mediating exercise-induced lipolysis. Muscle-derived IL-6 may also work to inhibit the effects of pro-inflammatory cytokines such as tumour necrosis factor α . The latter cytokine is produced by adipose tissue and inflammatory cells and appears to play a pathogenetic role in insulin resistance and atherogenesis.

Recent studies show that several cytokines can be detected in plasma during and after strenuous exercise (Pedersen & Hoffman-Goetz, 2000; Pedersen *et al.* 2001). Thus, the increase in tumour necrosis factor (TNF) α and interleukin (IL)-1 β levels is accompanied by a dramatic increase in IL-6. This increase is followed by increases in cytokine inhibitors (IL-1 receptor antagonist (IL-1ra), TNF receptors and the anti-inflammatory cytokine IL-10 (Ostrowski *et al.* 1998*a, b*, 1999). Also, concentrations of the chemokines, IL-8, macrophage inflammatory protein (MIP)-1 α and MIP-1 β , are elevated after a marathon (Ostrowski *et al.* 2001). In relation to exercise, IL-6 is produced in larger amounts than any other cytokine examined. Furthermore, it has been demonstrated that contracting skeletal muscles produce IL-6. The present review focuses on mechanisms underlying muscle-derived IL-6 and discusses for the first time possible biological roles of muscle-derived IL-6.

Characteristics of IL-6

IL-6 is a variably glycosylated protein with a molecular mass of 22–27 kDa, depending on the cellular source and the amount of post-translational modification. It is synthesized as a precursor protein of 212 amino acids, with a 28 amino acid signal sequence and a 184 amino acid mature segment (Haegeman *et al.* 1986; Hirano *et al.* 1986; May *et al.* 1986; Zilberstein *et al.* 1986). IL-6 is a member of a family of cytokines that includes leukaemia inhibitory factor, IL-11, ciliary neurotropic factor (CNTF), oncostatin M (OSM) and cardiotrophin-1 (CT-1) (Gadient & Patterson, 1999). Membership is based on similarities in helical protein structure and a shared receptor subunit (transmembrane glycoprotein (gp) 130) (Hirano *et al.* 1994; Hibi *et al.* 1996).

IL-6 has been classified as both a pro- and an anti-inflammatory cytokine, but the current view is that IL-6 has primarily anti-inflammatory effects. Infusion of IL-6 into humans results in fever but does not cause shock or capillary-leakage-like syndrome, as observed with the prototypical pro-inflammatory cytokines, IL-1 and TNF α (Rehman *et al.* 1997). Unlike IL-1 and TNF α , IL-6 does not upregulate major inflammatory mediators such as nitric oxide or matrix metalloproteinases. Rather, IL-6 appears to be the primary inducer of the hepatocyte-derived acute-phase proteins, many of which have anti-inflammatory properties. IL-6 directly inhibits the expression of TNF α and IL-1 and, furthermore, is a potent inducer of IL-1ra, which exerts anti-inflammatory activity by blocking IL-1 receptors, thereby preventing signal transduction of the pro-inflammatory IL-1 (Rehman *et al.* 1997).

IL-6 is produced by many different cell types, but the main sources *in vivo* are stimulated monocytes/macrophages, fibroblasts and vascular endothelial cells (Akira *et al.* 1993). Recently, however, much attention has been given to the striking observation that, during resting conditions, 10–35 % of the body's IL-6 is produced by adipose tissue (Mohamed-Ali *et al.* 1997).

Other cells known to express IL-6 include keratinocytes, osteoblasts, T cells, B cells, neutrophils, eosinophils, mast cells, smooth muscle cells and skeletal muscle cells (Akira *et al.* 1993). Typical stimuli for IL-6 production include IL-1, TNF α and bacterial endotoxin (Akira *et al.* 1993). Hypoxia induces IL-6 in cultured endothelial cells (Yan *et al.* 1995) and hypoxia *in vivo* elevates plasma IL-6 (Klausen *et al.* 1997).

IL-6 appears to play a central role in lipopolysaccharide (LPS)-induced fever (Sundgren-Andersson *et al.* 1998) and has growth-regulatory effects on a number of cells. Thus, IL-6 induces B and T cell differentiation and is involved in bone metabolism (Muraguchi *et al.* 1988; Houssiau *et al.* 1988; Takai *et al.* 1988).

Muscle damage and IL-6

Initially, it was thought that the cytokine response to exercise represented a reaction to exercise-induced muscle injury. Thus, we found that peak IL-6 concentrations were associated with prolonged muscle damage, using an eccentric exercise model in which the creatine kinase (CK) level peaked at day 4 after the exercise (Bruunsgaard *et al.* 1997). However, later studies from our group using exercise models in which CK peaked 1 day after exercise failed to show an association between peak IL-6 and peak CK levels (Ostrowski *et al.* 1998a, 1999). Furthermore, we have demonstrated recently, using an eccentric exercise model, that CK levels increased by up to 1000-fold with only a 4-fold increase in plasma IL-6 during the subsequent days (A. D. Toft, L. B. Jensen, H. Bruunsgaard, T. Ibfeldt, J. Halkjaer-Kristensen, M. A. Febbraio & B. K. Pedersen, unpublished data). The latter findings suggest that the huge increase in plasma levels of IL-6 in exercise models where the CK level does not change or is enhanced by a fewfold only, is related to mechanisms other than muscle damage. Also, a recent study (Croisier *et al.* 1999) failed to find an association between the increase in IL-6 and biochemical markers for muscle damage. These authors (Croisier *et al.* 1999) showed that training reduced the increase in myoglobin and decreased delayed-onset muscle soreness in response to a bout of eccentric exercise, whereas the increase in IL-6 was not influenced by training. It is most likely that the huge and immediate increase in IL-6 in response to long-duration exercise is independent of muscle damage, while muscle damage itself is followed by repair mechanisms including invasion of macrophages into the muscle leading to IL-6 production. IL-6 production in relation to muscle damage occurs later and is of smaller magnitude than IL-6 production related to muscle contractions. In a recent study, we tested the possibility that elevated levels of IL-6 during eccentric exercise might be inhibited by anti-oxidant treatment with vitamin C and vitamin E, which has been suggested to protect against muscle damage. However, we were unable to demonstrate any effect of anti-oxidants on plasma cytokines or muscle damage (Petersen *et al.* 2001). Non-steroid anti-inflammatory drugs might be used in future studies to evaluate the effect on IL-6, when the local inflammatory response is limited.

Kinetics of the IL-6 response to exercise

The finding of markedly increased levels of IL-6 after strenuous exercise has consistently been observed in many studies (Northoff & Berg, 1991; Sprenger *et al.* 1992; Ullum *et al.* 1994; Drenth *et al.* 1995; Nehlsen-

Cannarella *et al.* 1997; Castell *et al.* 1997; Rohde *et al.* 1997; Hellsten *et al.* 1997; Ostrowski *et al.* 1998a, b, 1999; Toft *et al.* 2000). A 2-fold increase in plasma IL-6 was demonstrated after 6 min of intense exercise (Nielsen *et al.* 1996). In treadmill running, the IL-6 level in blood was significantly enhanced 30 min after the start, with peak levels being attained after 2.5 h of running (Ostrowski *et al.* 1998a). In other studies, in which IL-6 was not measured during the running exercise but at several time points afterwards, maximal IL-6 levels were found immediately after the exercise, followed by a rapid decline. Thus, following a marathon run maximal IL-6 levels (100-fold increase) were measured immediately after the 3–3.5 h race (Ostrowski *et al.* 1998b, 1999).

In contrast, using a prolonged eccentric one-legged exercise model lasting 1 h (Rohde *et al.* 1997) or a two-legged high-intensity eccentric exercise model lasting 30 min (Bruunsgaard *et al.* 1997), the IL-6 level did not peak until 1–1.5 h after the exercise. In another study, subjects performed five bouts of one-legged eccentric exercise. The concentration of plasma IL-6 peaked 90 min after the exercise and was elevated for 4 days (Hellsten *et al.* 1997). It is very clear that the kinetics of the IL-6 response differ between that induced by concentric muscle contractions and that induced by eccentric exercise associated with muscle damage (Bruunsgaard *et al.* 1997; Rohde *et al.* 1997; Ostrowski *et al.* 1998b, 1999). In relation to concentric exercise, the increase in IL-6 is tightly related to the duration of the exercise (Ostrowski *et al.* 1998a), and there appears to be a logarithmic relationship between the increase in IL-6 and the duration of the exercise. IL-6 levels decline after the concentric exercise to reach pre-exercise values within a few hours. In contrast, eccentric exercise induces only a modest increase in plasma IL-6 and the IL-6 level peaks some time after cessation of the exercise and is elevated for several days.

Intensity of exercise and IL-6

Data from the Copenhagen Marathon race (1996, 1997 and 1998, $n = 56$) suggest that there is a correlation between the intensity of exercise and the increase in plasma IL-6 (Ostrowski *et al.* 2001). Furthermore, a correlation between peak IL-6 and heart rate was demonstrated (Ostrowski *et al.* 1998a). Results from an animal study suggested that the increase in adrenaline during stress was responsible for the increase in IL-6 (DeRijk *et al.* 1994). However, recent data from our group showed that when adrenaline was infused into volunteers to closely mimic the increase in plasma adrenaline during 2.5 h of running exercise, plasma IL-6 increased only 4-fold during the infusion, but 30-fold during the exercise (Steensberg *et al.* 2001b). Thus, it seems that adrenaline only plays a minor role in the exercise-induced increase in plasma IL-6. It has been demonstrated previously that peak plasma IL-6 during exercise correlated with plasma lactate (Ostrowski *et al.* 1998a). However, in a recent study

(Steensberg *et al.* 2001c), patients with mitochondrial myopathy, characterized by high plasma levels of lactate, were examined. They were treated with dichloroacetate (DCA) for 15 days. DCA inhibits pyruvate dehydrogenase kinase, thereby increasing the activity of the pyruvate dehydrogenase complex. The same exercise test was repeated on days with and without treatment. DCA lowered the plasma lactate and increased plasma IL-6 at rest. Furthermore, IL-6 increased in response to exercise only during DCA treatment. Thus, IL-6 production is not a direct result of high lactate levels.

Muscle-derived IL-6

A study was performed to test the hypothesis that inflammatory cytokines are produced in skeletal muscle in response to intense long-duration exercise (Ostrowski *et al.* 1998b). Muscle biopsies and blood samples were taken before and after a marathon race. The levels of IL-6 and IL-1ra proteins were markedly increased after the exercise. The plasma levels of IL-6 decreased, whereas those of IL-1ra increased further, 2 h after the exercise. A comparative polymerase chain reaction (PCR) technique was established to detect mRNA for cytokines in skeletal muscle biopsies and blood mononuclear cells (BMNCs). Before exercise, IL-6 mRNA could not be detected in muscle or BMNCs. In contrast, IL-6 mRNA was detected in muscle biopsies after exercise but not in the BMNC samples.

Starkie *et al.* (2000, 2001b) observed that the intracellular levels of IL-6 and IL-1ra protein in circulating monocytes were not augmented in response to exercise, and concluded that these cells did not contribute to the exercise-induced increases in plasma IL-6. The finding that blood cells do not contribute to the exercise-induced increase in IL-1 β and IL-6 plasma levels was further substantiated using a quantitative PCR approach. Quantification of the IL-1 β and IL-6 mRNA levels in BMNCs during exercise demonstrated no change despite an increase in the plasma level of the proteins (Moldoveanu *et al.* 2000). Recently, we measured IL-1 β and IL-6 mRNA levels in muscle biopsies using quantitative competitive RT-PCR and found that both specific mRNAs increased more than 10-fold following a marathon (Schjerling *et al.* 2001). The finding of IL-6 mRNA in muscle in response to exercise was confirmed in a rat exercise model, using the quantitative competitive RT-PCR method (Jonsdottir *et al.* 2000). In this model, rats were subjected to electrically stimulated eccentric or concentric contractions of one hind leg, while the other leg remained at rest. The eccentric and concentric contractions both resulted in elevated levels of IL-6 mRNA in the exercised muscle, whereas the level in the resting leg was not elevated. The finding of similar levels of IL-6 mRNA in both concentric- and eccentric-exercised muscle indicates that the cytokine production cannot be as closely related to muscle damage as was first thought. It does, however, appear that the local IL-6 production is

connected with exercising muscle – and is not due to a systemic effect – because IL-6 mRNA was elevated only in the muscle from the exercised leg, and not in the other resting leg.

Recently, we demonstrated that the net IL-6 release from contracting skeletal muscles could more than account for the exercise-induced increase in arterial plasma concentrations (Steensberg *et al.* 2000). Moreover, by obtaining arterial-femoral venous differences from an exercising leg, we found that exercising muscles released IL-6. In addition, during the last 2 h of exercise the release per unit time was ~ 17 -fold higher than the amount accumulating in the plasma. Although IL-6 appears to be produced in the muscle, which cell types within the muscle are responsible for the production remains unknown. Myoblasts have been shown to be able to produce IL-6 (Bartoccioni *et al.* 1994; De Rossi *et al.* 2000), but endothelial cells (Yan *et al.* 1995), fibroblasts (De Rossi *et al.* 2000) and smooth muscle (Klouché *et al.* 1999) have also been shown to produce IL-6 under certain conditions. *In situ* hybridization for IL-6 mRNA in mouse hearts after LPS stimulation gave a low signal from cardiomyocytes but a greater signal from non-myocyte cells (Saito *et al.* 2000). *In situ* hybridization on damaged muscles showed increased expression of IL-6 mRNA inside muscle fibres as well as in mononucleated cells (Kami & Senba, 1998).

Carbohydrate ingestion and IL-6

Several studies have reported that carbohydrate ingestion attenuates elevations in plasma IL-6 during both running and cycling (Nehlsen-Cannarella *et al.* 1997; Nieman *et al.* 1998). In contrast, researchers from Melbourne, Australia (Starkie *et al.* 2000) reported that plasma IL-6 was unaffected by carbohydrate ingestion during cycling. However, in that study the subjects were highly endurance trained and plasma IL-6 increased to only ~ 2 pg ml $^{-1}$, even without carbohydrate ingestion. This increase is markedly less than that previously observed in moderately trained subjects (Nieman *et al.* 1998). Recently, the same group reported that carbohydrate ingestion did attenuate the increase in plasma IL-6 found in response to both cycling and running (Starkie *et al.* 2001a). In the latter experiment, subjects of similar aerobic fitness to those previously reported were used (Nieman *et al.* 1998). Furthermore, IL-6 gene expression in the muscles was investigated. However, the level of IL-6 mRNA was found not to be affected by carbohydrate ingestion. IL-6 protein release from contracting muscle was not measured, and it is possible that carbohydrate ingestion did affect the release of IL-6, without altering the gene expression.

These authors suggested that it is the liver rather than contracting muscles that is responsible for the differences in plasma IL-6 between the carbohydrate and the non-carbohydrate trial. Moreover, it is possible that carbohydrate ingestion may have blunted IL-6

production in hepatic tissue. In addition, differences in the clearance of IL-6 may be responsible for the attenuated plasma IL-6 during carbohydrate ingestion. Further investigation is needed to clarify whether organs other than contracting muscles are involved in IL-6 production during exercise.

Muscle derived IL-6 may act as a hormone

During the past 30 years, there has been intensive research to unravel the mechanisms that regulate the release of glucose from the liver to the blood during physical exercise, so that the blood glucose level is maintained in spite of increased glucose uptake in working skeletal muscles (Kjaer, 2001). Research has demonstrated that exercise-induced changes in insulin and/or glucagons (Kjaer, 2001), cortisol (Cryer, 1993), adrenaline (Howlett *et al.* 1999) or adrenergic neural stimulation (Sigal *et al.* 1994, 2000) cannot, by themselves, account for the exercise-induced increase in hepatic glucose production. Indeed, it has been concluded that the possibility exists that an as yet unidentified factor, released from contracting muscle cells, may contribute to the increase in hepatic glucose production (Howlett *et al.* 1999).

An elevated plasma IL-6 response was found when subjects exercised in a glycogen-depleted state (Gleeson & Bishop, 2000). Recently, we presented data showing that release of IL-6 from a glycogen-depleted exercising leg occurred 1 h before release was observed in a non-depleted control leg (Keller *et al.* 2001; Steensberg *et al.* 2001a). Furthermore, when individual data with regard to the legs with the lowest glycogen content at the end of exercise were analysed, it was demonstrated that the highest levels of IL-6 mRNA were expressed in these legs. Therefore, the glycogen content or an energy crisis in the contracting muscle may be one stimulus for the IL-6 release. IL-6 has been shown to have a marked influence on hepatic glucose metabolism. Moreover, IL-6 has also been shown to inhibit glycogen synthase activity and accelerate glycogen phosphorylase activity (Kanemaki *et al.* 1998). In addition, it has been demonstrated that injection of recombinant human IL-6 (rhIL-6) into humans increases hepatic glucose production (Stouthard *et al.* 1995) and the fasting blood glucose concentration in a dose-dependent manner (Tsigos *et al.* 1997). These data raise the possibility that the IL-6 produced by contracting skeletal muscle partly mediates the hepatic glucose output necessary to maintain blood glucose homeostasis when the uptake of blood glucose by skeletal muscles is increased by prolonged exercise (Fig. 1).

Besides the glucoregulatory effect of IL-6, this cytokine may also be involved in other metabolic pathways (Fig. 1). Concomitantly with the increase in liver glucose output during IL-6 infusion, Stouthard *et al.* (1996) observed an increased release of free fatty acids (FFAs). Furthermore, infusion of IL-6 into rats increased serum triglyceride and FFA levels in a dose-dependent manner (Nonogaki *et al.* 1995). The hypertriglyceridaemia was

due to increased secretion by the liver and not to decreased clearance. The increased levels of serum FFAs were due to lipolysis.

Increased glucose transport was found in jejunal tissue incubated with IL-6, compared with controls (Hardin *et al.* 2000). Moreover, IL-6 also appears to be able to increase the absorption of glucose in the gut, thereby increasing the plasma glucose levels. Furthermore, there is evidence to suggest that IL-6 may augment glucose uptake by insulin-sensitive tissues. Stouthard *et al.* (1996) demonstrated that infusion of rhIL-6 into human subjects increased whole body glucose disposal and subsequent oxidation compared with a control trial. Even though endogenous glucose production was increased with rhIL-6 infusion, the metabolic clearance rate of glucose was higher in this trial, suggesting that relative hyperglycaemia was not responsible for the augmented glucose disposal. Moreover, in a follow-up study, it was demonstrated that IL-6 increased both basal and insulin-stimulated glucose uptake in cultured 3T3-L1 adipocytes (Stouthard *et al.* 1996). These authors concluded that IL-6 acted by increasing glucose transporter (GLUT)-1, intrinsic activity. Although highly speculative, it is possible that IL-6 produced by the muscle may act on signalling molecules, which enhance translocation of GLUT-4, the transporter responsible for facilitating contraction-mediated glucose uptake.

It is tempting to speculate that IL-6 may be involved in glucoregulatory processes in circumstances in which glucose homeostasis is disrupted. Basal serum IL-6 is higher in type II diabetics (Pickup *et al.* 2000) and in cancer patients with insulin resistance (Makino *et al.* 1998), while a polymorphism of the IL-6 gene influences insulin sensitivity in humans (Fernandez-Real *et al.* 2000). Interestingly, in the study by Pickup *et al.* (2000) the basal production of IL-6 in cultured blood cells from diabetics was depressed compared with blood from non-diabetics. Therefore, the possibility exists that the elevated serum IL-6 is a consequence of an increased production and release of IL-6 from insulin-sensitive tissue in response to the impaired insulin sensitivity. Based on this information, it is of utmost interest to investigate the role of muscle-derived IL-6 for metabolic homeostasis, including substrate uptake and release in different tissues.

Does IL-6 have a local effect on the muscle tissue?

Another role for the exercise-produced IL-6 may be to regulate processes within the producing muscles (Fig. 1). IL-6 is known to have local regulatory functions in other tissues such as the nervous system (Van Wagoner & Benveniste, 1999) and the skeleton (Franchimont *et al.* 1997).

The downstream signalling pathway for IL-6 has been well investigated, especially in liver cells. IL-6 belongs to a family of cytokines which bind to different specific

receptors but which all transmit signals over the cell membrane via the gp130 tyrosine kinase receptor. IL-6 binds to the IL-6 receptor on the surface of target cells, which then causes dimerization of the tyrosine kinase receptor gp130. Upon dimerization, gp130 recruits cytoplasmic Janus kinases (JAKs), which then phosphorylate signal transducer and activator of transcription 3 (STAT3). The phosphorylated STAT3 then dimerize and translocate from the cytoplasm into the nucleus where STAT3 activates transcription of a number of different genes. Besides the more specific signal transduction pathway via STAT3, the dimerization of gp130 also causes activation of the phosphatidylinositol-3 kinase (PI-3 kinase) and the mitogen-activated protein kinase (MAPK) cascade. An important review on the gp130 signal transduction pathway is given by Hirano *et al.* (1997). gp130 is ubiquitously expressed throughout the body (Hibi *et al.* 1990; Saito *et al.* 1992), whereas the distribution of the IL-6 receptor is more restricted. IL-6 receptor mRNA is found in skeletal muscle, although at low levels, but the level is subject to inducing signals as demonstrated by LPS stimulation (Zhang *et al.* 2000). However, muscles do not necessarily need to express the IL-6 receptor, since serum contains a soluble form of the receptor which can substitute for the membrane-bound

receptor (Croucher *et al.* 1999). What then could be the possible function of a local IL-6 signal? Myoblast proliferation is stimulated by IL-6 and satellite cell proliferation is regulated by an autocrine secretion of IL-6 (Austin & Burgess 1991; Austin *et al.* 1992; Cantini *et al.* 1995). Furthermore, IL-6 administration also enhances differentiation of the myoblasts (Okazaki *et al.* 1996). This has led to the idea that IL-6 may be involved in hypertrophy during resistance exercise (Vierck *et al.* 2000). However, IL-6 production from the muscles is mainly seen after endurance exercise rather than after resistance exercise (Ostrowski *et al.* 2000). Overexpression of IL-6 in transgenic mice causes muscle atrophy and increased levels of cathepsins in muscle, indicating that IL-6 is involved in regulating muscle protein breakdown (Tsujinaka *et al.* 1995). This atrophy was completely abolished following injection of antibodies against the IL-6 receptor (Tsujinaka *et al.* 1996). However, this effect on muscles is not necessarily a direct effect, since for instance the circulating level of insulin-like growth factor 1 (IGF-1) also decreased in the IL-6 transgenic mice (De Benedetti *et al.* 1997). IL-6 can induce the expression of vascular endothelial growth factor (VEGF) (Cohen *et al.* 1996) and could therefore be responsible for the upregulation of VEGF seen after exercise (Gustafsson *et*




Figure 1

Schematic presentation of the possible biological effects of muscle-derived IL-6. TG, triglyceride; FFA, free fatty acid.

al. 1999), suggesting a role in capillary growth. Since systemic IL-6 has well-known effects on glucose metabolism in the liver and fatty acid metabolism in the adipose tissue as described above, another potential function of local IL-6 may well be to regulate the metabolism within the muscle by exerting similar or perhaps opposite effects on the muscle tissue. The actual local role of exercise-produced IL-6 in muscle has still to be determined.

How does muscle-derived IL-6 interact with IL-6 produced by inflammatory cells or adipose tissue?

Given the many beneficial effects of physical exercise on health, it is hard to believe that muscle-derived IL-6 is detrimental to health. However, elevated levels of circulating IL-6 have been associated with several disorders. Thus, increased levels of TNF and IL-6 have been observed in obese individuals, in smokers and in non-insulin-dependent diabetes mellitus (Vgontzas *et al.* 2000). In several population-based studies, plasma concentrations of IL-6 have been shown to predict total and cardiovascular mortality (Ridker *et al.* 2000a, b). Furthermore, ageing is associated with increased levels of both TNF α and IL-6 (Bruunsgaard *et al.* 1999). It has been proposed that IL-6 is the mediator that links the acute phase response to visceral obesity, insulin resistance and atherosclerosis (Yudkin *et al.* 2000; and Dr Robert S. Munford's Grand Rounds of July 6, 2000).

High levels of IL-6 in patients with metabolic syndrome may be explained by the fact that IL-6 is produced in adipose tissue (Mohamed-Ali *et al.* 1997; Fried *et al.* 1998). Omental adipose tissue produces more IL-6 than subcutaneous tissue (Fried *et al.* 1998). Adipose tissue also produces and releases TNF α (Tsigos *et al.* 1999), and TNF α is thought to play a mechanistic role in insulin resistance. Thus, TNF α downregulates GLUT-4 and inhibits insulin receptor activity (Hotamisligil, 1999). Since TNF α can trigger IL-6 release, one theory holds that it is adipose tissue TNF α that is actually the 'driver' behind the metabolic syndrome. In support of this theory is the finding that in elderly people serum levels of leptin and TNF α were correlated even after the effect of sex and body mass index was taken into account (Bruunsgaard *et al.* 2000a, b). It was also demonstrated that blood levels of soluble TNF α receptors (an indicator of TNF activation) correlate with blood leptin levels in both controls and diabetic subjects (Mantzoros *et al.* 1997). Furthermore, TNF α was elevated in elderly patients with atherosclerosis compared with age-matched subjects without this diagnosis (Bruunsgaard *et al.* 1999, 2000a, b). However, although TNF α correlated with IL-6 (Bruunsgaard *et al.* 1999), the latter cytokine was not associated with atherosclerosis. In addition, TNF α is the prototype of a pro-inflammatory cytokine, whereas IL-6 is now regarded as an anti-inflammatory cytokine. Thus, it is possible that TNF α , rather than IL-6, should be regarded as the cytokine that induces insulin resistance and

thereby initiates diabetes type 2 and atherosclerosis. According to this theory, muscle-derived IL-6 may work to inhibit the effects of TNF α and IL-6 may represent an important messenger from muscle work to health effects (Fig. 1). An alternative explanation is that muscle-derived IL-6 represents an isoform that differs from the IL-6 molecules produced by adipose tissue and inflammatory cells.

Conclusion

Muscle-derived IL-6 is released into the circulation in large amounts and is likely to work in a hormone-like fashion, exerting its effect on the liver and adipose tissue, thereby contributing to the maintenance of glucose homeostasis during exercise and mediating exercise-induced lipolysis. Muscle-derived IL-6 may also work to inhibit the effects of pro-inflammatory cytokines such as TNF α , thereby protecting against insulin resistance and atherogenesis.

- AKIRA, S., TAGA, T. & KISHIMOTO, T. (1993). Interleukin-6 in biology and medicine. *Advances in Immunology* **54**, 1–78.
- AUSTIN, L., BOWER, J., KUREK, J. & VAKAKIS, N. (1992). Effects of leukaemia inhibitory factor and other cytokines on murine and human myoblast proliferation. *Journal of the Neurological Sciences* **112**, 185–191.
- AUSTIN, L. & BURGESS, A. W. (1991). Stimulation of myoblast proliferation in culture by leukaemia inhibitory factor and other cytokines. *Journal of the Neurological Sciences* **101**, 193–197.
- BARTOCCIONI, E., MICHAELIS, D. & HOHLFELD, R. (1994). Constitutive and cytokine-induced production of interleukin-6 by human myoblasts. *Immunology Letters* **42**, 135–138.
- BRUUNSGAARD, H., ANDERSEN-RANBERG, K., JEUNE, B., PEDERSEN, A. N., SKINHOJ, P. & PEDERSEN, B. K. (1999). A high plasma concentration of TNF-alpha is associated with dementia in centenarians. *Journals of Gerontology* **54A**, 357–364.
- BRUUNSGAARD, H., GALBO, H., HALKJAER-KRISTENSEN, J., JOHANSEN, T. L., MACLEAN, D. A. & PEDERSEN, B. K. (1997). Exercise-induced increase in serum interleukin-6 in humans is related to muscle damage. *Journal of Physiology* **499**, 833–841.
- BRUUNSGAARD, H., PEDERSEN, A. N., SCHROLL, M., SKINHOJ, P. & PEDERSEN, B. K. (2000a). TNF- α , leptin, and lymphocyte function in human aging. *Life Sciences* **67**, 2721–2731.
- BRUUNSGAARD, H., SKINHOJ, P., PEDERSEN, A. N., SCHROLL, M. & PEDERSEN, B. K. (2000b). Ageing, tumour necrosis factor- α (TNF- α) and atherosclerosis. *Clinical and Experimental Immunology* **121**, 255–260.
- CANTINI, M., MASSIMINO, M. L., RAPIZZI, E., ROSSINI, K., CATANI, C., DALLA, L. L. & CARRARO, U. (1995). Human satellite cell proliferation in vitro is regulated by autocrine secretion of IL-6 stimulated by a soluble factor(s) released by activated monocytes. *Biochemical and Biophysical Research Communications* **216**, 49–53.
- CASTELL, L. M., POORTMANS, J. R., LECLERCQ, R., BRASSEUR, M., DUCHATEAU, J. & NEWSHOLME, E. A. (1997). Some aspects of the acute phase response after a marathon race, and the effects of glutamine supplementation. *European Journal of Applied Physiology and Occupational Physiology* **75**, 47–53.

- COHEN, T., NAHARI, D., CEREM, L. W., NEUFELD, G. & LEVI, B. Z. (1996). Interleukin 6 induces the expression of vascular endothelial growth factor. *Journal of Biological Chemistry* **271**, 736–741.
- CROISIER, J. L., CAMUS, G., VENNEMAN, I., DEBY-DUPONT, G., JUCHMES-FERIR, A., LAMY, M., CRIELAARD, J. M., DEBY, C. & DUCHATEAU, J. (1999). Effects of training on exercise-induced muscle damage and interleukin 6 production. *Muscle and Nerve* **22**, 208–212.
- CROUCHER, P. I., WANG, F. & HARGREAVES, P. G. (1999). Interleukin-6 receptor shedding: a possible role for members of the ADAM family. *Biochemical Society Transactions* **27**, 224–228.
- CRYER, P. E. (1993). Glucose counterregulation: prevention and correction of hypoglycemia in humans. *American Journal of Physiology* **264**, E149–155.
- DE BENEDETTI, F., ALONZI, T., MORETTA, A., LAZZARO, D., COSTA, P., POLI, V., MARTINI, A., CILIBERTO, G. & FATTORI, E. (1997). Interleukin 6 causes growth impairment in transgenic mice through a decrease in insulin-like growth factor-I. A model for stunted growth in children with chronic inflammation. *Journal of Clinical Investigation* **99**, 643–650.
- DERLIJK, R. H., BOELEN, A., TILDERS, F. J. & BERKENBOSCH, F. (1994). Induction of plasma interleukin-6 by circulating adrenaline in the rat. *Psychoneuroendocrinology* **19**, 155–163.
- DE ROSSI, M., BERNASCONI, P., BAGGI, F., DE WAAL, M. R. & MANTEGAZZA, R. (2000). Cytokines and chemokines are both expressed by human myoblasts: possible relevance for the immune pathogenesis of muscle inflammation. *International Immunology* **12**, 1329–1335.
- DRENTH, J. P., VAN UUM, S. H., VAN DEUREN, M., PESMAN, G. J., VAN DER VEN JONGEKRUIG, J. & VAN DER MEER, J. W. (1995). Endurance run increases circulating IL-6 and IL-1ra but downregulates ex vivo TNF- α and IL-1 β production. *Journal of Applied Physiology* **79**, 1497–1503.
- FERNANDEZ-REAL, J. M., BROCH, M., VENDRELL, J., GUTIERREZ, C., CASAMITJANA, R., PUJEAT, M., RICHART, C. & RICART, W. (2000). Interleukin-6 gene polymorphism and insulin sensitivity. *Diabetes* **49**, 517–520.
- FRANCHIMONT, N., GANGJI, V., DURANT, D. & CANALIS, E. (1997). Interleukin-6 with its soluble receptor enhances the expression of insulin-like growth factor-I in osteoblasts. *Endocrinology* **138**, 5248–5255.
- FRIED, S. K., BUNKIN, D. A. & GREENBERG, A. S. (1998). Omental and subcutaneous adipose tissues of obese subjects release interleukin-6: depot difference and regulation by glucocorticoid. *Journal of Clinical Endocrinology and Metabolism* **83**, 847–850.
- GADIENT, R. A. & PATTERSON, P. H. (1999). Leukemia inhibitory factor, interleukin 6, and other cytokines using the GP130 transducing receptor: roles in inflammation and injury. *Stem Cells* **17**, 127–137.
- GLEESON, M. & BISHOP, N. C. (2000). Special feature for the Olympics: effects of exercise on the immune system: modification of immune responses to exercise by carbohydrate, glutamine and anti-oxidant supplements. *Immunology and Cell Biology* **78**, 554–561.
- GUSTAFSSON, T., PUNTSCHART, A., KAIJSER, L., JANSSON, E. & SUNDBERG, C. J. (1999). Exercise-induced expression of angiogenesis-related transcription and growth factors in human skeletal muscle. *American Journal of Physiology* **276**, H679–685.
- HAEGEMAN, G., CONTENT, J., VOLCKAERT, G., DERYNCK, R., TAVERNIER, J. & FIERIS, W. (1986). Structural analysis of the sequence coding for an inducible 26-kDa protein in human fibroblasts. *European Journal of Biochemistry* **159**, 625–632.
- HARDIN, J., KROEGER, K., CHUNG, B. & GALL, D. G. (2000). Effect of proinflammatory interleukins on jejunal nutrient transport. *Gut* **47**, 184–191.
- HELLSTEN, Y., FRANDSEN, U., ØRTHENBLAD, N., SJÖDIN, N. & RICHTER, E. A. (1997). Xanthine oxidase in human skeletal muscle following eccentric exercise: a role in inflammation. *Journal of Physiology* **498**, 239–248.
- HIBI, M., MURAKAMI, M., SAITO, M., HIRANO, T., TAGA, T. & KISHIMOTO, T. (1990). Molecular cloning and expression of an IL-6 signal transducer, gp 130. *Cell* **63**, 1149–1157.
- HIBI, M., NAKAJIMA, K. & HIRANO, T. (1996). IL-6 cytokine family and signal transduction: a model of the cytokine system. *Journal of Molecular Medicine* **74**, 1–12.
- HIRANO, T., MATSUDA, T. & NAKAJIMA, K. (1994). Signal transduction through gp130 that is shared among the receptors for the interleukin 6 related cytokine subfamily. *Stem Cells* **12**, 262–277.
- HIRANO, T., NAKAJIMA, K. & HIBI, M. (1997). Signaling mechanisms through gp130: a model of the cytokine system. *Cytokine and Growth Factor Reviews* **8**, 241–252.
- HIRANO, T., YASUKAWA, K., HARADA, H., TAGA, T., WATANABE, Y., MATSUDA, T., KASHIWAMURA, S., NAKAJIMA, K., KOYAMA, K., IWAMATSU, A. *et al.* (1986). Complementary DNA for a novel human interleukin (BSF-2) that induces B lymphocytes to produce immunoglobulin. *Nature* **324**, 73–76.
- HOTAMISLIGIL, G. S. (1999). The role of TNF α and TNF receptors in obesity and insulin resistance. *Journal of Internal Medicine* **245**, 621–625.
- HOUSIAU, F. A., COULIE, P. G., OLIVE, D. & VAN SNICK, J. (1988). Synergistic activation of human T cells by interleukin 1 and interleukin 6. *European Journal of Immunology* **18**, 653–656.
- HOWLETT, K., FEBBRAIO, M. & HARGREAVES, M. (1999). Glucose production during strenuous exercise in humans: role of epinephrine. *American Journal of Physiology* **276**, E1130–1135.
- JONSDOTTIR, I. H., SCHJERLING, P., OSTROWSKI, K., ASP, S., RICHTER, E. A. & PEDERSEN, B. K. (2000). Muscle contractions induce interleukin-6 mRNA production in rat skeletal muscles. *Journal of Physiology* **528**, 157–163.
- KAMI, K. & SENBA, E. (1998). Localization of leukemia inhibitory factor and interleukin-6 messenger ribonucleic acids in regenerating rat skeletal muscle. *Muscle and Nerve* **21**, 819–822.
- KANEMAKI, T., KITADE, H., KAIBORI, M., SAKITANI, K., HIRAMATSU, Y., KAMIYAMA, Y., ITO, S. & OKUMURA, T. (1998). Interleukin 1 β and interleukin 6, but not tumor necrosis factor α , inhibit insulin-stimulated glycogen synthesis in rat hepatocytes. *Hepatology* **27**, 1296–1303.
- KELLER, C., STEENSBERG, A., PILEGAARD, H., OSADA, T., SALTIN, B., PEDERSEN, B. K. & NEUFER, P. D. (2001). Transcriptional activation of the IL-6 gene in human contracting skeletal muscle – influence of muscle glycogen content. *FASEB Journal* (in the Press).
- KJAER, M. (2001). Hepatic fuel metabolism during exercise. In *Exercise Metabolism*, ed. HARGREAVES, M., pp. 73–98. Human Kinetics, Champaign, IL, USA.
- KLAUSEN, T., OLSEN, N. V., POULSEN, T. D., RICHALET, J. P. & PEDERSEN, B. K. (1997). Hypoxemia increases serum interleukin-6 in humans. *European Journal of Applied Physiology and Occupational Physiology* **76**, 480–482.
- KLOUCHE, M., BHAKDI, S., HEMMES, M. & ROSE-JOHN, S. (1999). Novel path to activation of vascular smooth muscle cells: up-regulation of gp130 creates an autocrine activation loop by IL-6 and its soluble receptor. *Journal of Immunology* **163**, 4583–4589.

- MAKINO, T., NOGUCHI, Y., YOSHIKAWA, T., DOI, C. & NOMURA, K. (1998). Circulating interleukin 6 concentrations and insulin resistance in patients with cancer. *British Journal of Surgery* **85**, 1658–1662.
- MANTZOROS, C. S., MOSCHOS, S., AVRAMOPOULOS, I., KAKLAMANI, V., LIOLIOS, A., DOULGERAKIS, D. E., GRIVEAS, I., KATSIAMBROS, N. & FLIER, J. S. (1997). Leptin concentrations in relation to body mass index and the tumor necrosis factor- α system in humans. *Journal of Clinical Endocrinology and Metabolism* **82**, 3408–3413.
- MAY, L. T., HELFGOTT, D. C. & SEHGAL, P. B. (1986). Anti- β -interferon antibodies inhibit the increased expression of HLA-B7 mRNA in tumor necrosis factor-treated human fibroblasts: structural studies of the β 2 interferon involved. *Proceedings of the National Academy of Sciences of the USA* **83**, 8957–8961.
- MOHAMED-ALI, V., GOODRICK, S., RAWESH, A., KATZ, D. R., MILES, J. M., YUDKIN, J. S., KLEIN, S. & COPPACK, C. (1997). Subcutaneous adipose tissue releases interleukin-6, but not tumour necrosis factor- α , in vivo. *Journal of Clinical Endocrinology and Metabolism* **82**, 4196–4200.
- MOLDOVEANU, A. I., SHEPHARD, R. J. & SHEK, P. N. (2000). Exercise elevates plasma levels but not gene expression of IL-1 β , IL-6, and TNF- α in blood mononuclear cells. *Journal of Applied Physiology* **89**, 1499–1504.
- MURAGUCHI, A., HIRANO, T., TANG, B., MATSUDA, T., HORII, Y., NAKAJIMA, K. & KISHIMOTO, T. (1988). The essential role of B cell stimulatory factor 2 (BSF-2/IL-6) for the terminal differentiation of B cells. *Journal of Experimental Medicine* **167**, 332–344.
- NEHLSSEN-CANNARELLA, S. L., FAGOAGA, O. R., NIEMAN, D. C., HENSON, D. A., BUTTERWORTH, D. E., SCHMITT, R. L., BAILEY, E. M., WARREN, B. J., UTTER, A. & DAVIS, J. M. (1997). Carbohydrate and the cytokine response to 2.5 hours of running. *Journal of Applied Physiology* **82**, 1662–1667.
- NIELSEN, H. B., SECHER, N. H., CHRISTENSEN, N. J. & PEDERSEN, B. K. (1996). Lymphocytes and NK cell activity during repeated bouts of maximal exercise. *American Journal of Physiology* **271**, R222–227.
- NIEMAN, D. C., NEHLSSEN-CANNARELLA, S. L., FAGOAGA, O. R., HENSON, D. A., UTTER, A., DAVIS, J. M., WILLIAMS, F. & BUTTERWORTH, D. E. (1998). Influence of mode and carbohydrate on the cytokine response to heavy exertion. *Medicine and Science in Sports and Exercise* **30**, 671–678.
- NONOGAKI, K., FULLER, G. M., FUENTES, N. L., MOSER, A. H., STAPRANS, I., GRUNFELD, C. & FEINGOLD, K. R. (1995). Interleukin-6 stimulates hepatic triglyceride secretion in rats. *Endocrinology* **136**, 2143–2149.
- NORTHOFF, H. & BERG, A. (1991). Immunologic mediators as parameters of the reaction to strenuous exercise. *International Journal of Sports Medicine* **12**, S9–15.
- OKAZAKI, S., KAWAI, H., ARII, Y., YAMAGUCHI, H. & SAITO, S. (1996). Effects of calcitonin gene-related peptide and interleukin 6 on myoblast differentiation. *Cell Proliferation* **29**, 173–182.
- OSTROWSKI, K., HERMANN, C., BANGASH, A., SCHJERLING, P., NIELSEN, J. N. & PEDERSEN, B. K. (1998a). A trauma-like elevation in plasma cytokines in humans in response to treadmill running. *Journal of Physiology* **508**, 949–953.
- OSTROWSKI, K., ROHDE, T., ASP, S., SCHJERLING, P. & PEDERSEN, B. K. (1999). Pro- and anti-inflammatory cytokine balance in strenuous exercise in humans. *Journal of Physiology* **515**, 287–291.
- OSTROWSKI, K., ROHDE, T., ASP, S., SCHJERLING, P. & PEDERSEN, B. K. (2001). Chemokines are elevated in plasma after strenuous exercise. *European Journal of Applied Physiology* **84**, 244–245.
- OSTROWSKI, K., ROHDE, T., ZACHO, M., ASP, S. & PEDERSEN, B. K. (1998b). Evidence that IL-6 is produced in skeletal muscle during intense long-term muscle activity. *Journal of Physiology* **508**, 949–953.
- OSTROWSKI, K., SCHJERLING, P. & PEDERSEN, B. K. (2000). Physical activity and plasma interleukin-6 in humans – effect of intensity of exercise. *European Journal of Applied Physiology* **83**, 512–515.
- PEDERSEN, B. K. & HOFFMAN-GOETZ, L. (2000). Exercise and the immune system: regulation, integration and adaptation. *Physiological Reviews* **80**, 1055–1081.
- PEDERSEN, B. K., STEENBERG, A. & SCHJERLING, P. (2001). Exercise and interleukin-6. *Current Opinion in Hematology* **8**, 137–141.
- PETERSEN, E. W., OSTROWSKI, K., IBFELT, T., RICHELLE, M., OFFORD, E., HALKJAER-KRISTENSEN, J. & PEDERSEN, B. K. (2001). Effect of vitamin supplementation on the cytokine response and on muscle damage following strenuous exercise. *American Journal of Physiology* **280**, C1570–1575.
- PICKUP, J. C., CHUSNEY, G. D., THOMAS, S. M. & BURT, D. (2000). Plasma interleukin-6, tumour necrosis factor α and blood cytokine production in type 2 diabetes. *Life Sciences* **67**, 291–300.
- REHMAN, J., MILLS, P. J., CARTER, S. M., CHOU, J., THOMAS, J. & MAISEL, A. S. (1997). Dynamic exercise leads to an increase in circulating ICAM-1: further evidence for adrenergic modulation of cell adhesion. *Brain, Behavior, and Immunity* **11**, 343–351.
- RIDKER, P. M., HENNEKENS, C. H., BURING, J. E. & RIFAI, N. (2000a). C-reactive protein and other markers of inflammation in the prediction of cardiovascular disease in women. *New England Journal of Medicine* **342**, 836–843.
- RIDKER, P. M., RIFAI, N., STAMPFER, M. J. & HENNEKENS, C. H. (2000b). Plasma concentration of interleukin-6 and the risk of future myocardial infarction among apparently healthy men. *Circulation* **101**, 1767–1772.
- ROHDE, T., MACLEAN, D. A., RICHTER, E. A., KIENS, B. & PEDERSEN, B. K. (1997). Prolonged submaximal eccentric exercise is associated with increased levels of plasma IL-6. *American Journal of Physiology* **273**, E85–91.
- SAITO, H., PATTERSON, C., HU, Z., RUNGE, M. S., TIPNIS, U., SINHA, M. & PAPACONSTANTINOU, J. (2000). Expression and self-regulatory function of cardiac interleukin-6 during endotoxemia. *American Journal of Physiology – Heart and Circulatory Physiology* **279**, H2241–2248.
- SAITO, M., YOSHIDA, K., HIBI, M., TAGA, T. & KISHIMOTO, T. (1992). Molecular cloning of a murine IL-6 receptor-associated signal transducer, gp130, and its regulated expression in vivo. *Journal of Immunology* **148**, 4066–4071.
- SCHJERLING, P., OSTROWSKI, K., ASP, S. & PEDERSEN, B. K. (2001). Exercise induces interleukin-6 and interleukin-1 β expression within human skeletal muscle. *Journal of Physiology* **531**, P, 59–60P.
- SIGAL, R. J., FISHER, S. J., MANZON, A., MORAIS, J. A., HALTER, J. B., VRANIC, M. & MARLISS, E. B. (2000). Glucoregulation during and after intense exercise: effects of α -adrenergic blockade. *Metabolism* **49**, 386–394.
- SIGAL, R. J., PURDON, C., BILINSKI, D., VRANIC, M., HALTER, J. B. & MARLISS, E. B. (1994). Glucoregulation during and after intense exercise: effects of β -blockade. *Journal of Clinical Endocrinology and Metabolism* **78**, 359–366.
- SPRENGER, H., JACOBS, C., NAIN, M., GRESSNER, A. M., PRINZ, H., WESEMANN, W. & GEMSA, D. (1992). Enhanced release of cytokines, interleukin-2 receptors, and neopterin after long-distance running. *Clinical Immunology and Immunopathology* **63**, 188–195.

- STARKIE, R. L., ANGUS, D. J., ROLLAND, J., HARGREAVES, M. & FEBBRAIO, M. A. (2000). Effect of prolonged submaximal exercise and carbohydrate ingestion on monocyte intracellular cytokine production in humans. *Journal of Physiology* **528**, 647–655.
- STARKIE, R. L., ARKINSTALL, M. J., KOUKOULAS, I., HAWLEY, J. A. & FEBBRAIO, M. A. (2001*a*). Carbohydrate ingestion attenuates the increase in plasma interleukin-6, but not skeletal muscle interleukin-6 mRNA, during exercise in humans. *Journal of Physiology* **533**, 585–591.
- STARKIE, R. L., ROLLAND, J., ANGUS, D. J., AANDERSON, M. J. & FEBBRAIO, M. A. (2001*b*). Circulating monocytes are not the source of elevations in plasma IL-6 and TNF- α levels after prolonged running. *American Journal of Physiology – Cell Physiology* **280**, C769–774.
- STEENBERG, A., FEBBRAIO, M. A., OSADA, T., SCHJERLING, P., VAN HALL, G., SALTIN, B. & PEDERSEN, B. K. (2001*a*). Interleukin-6 production in contracting human skeletal muscle is influenced by pre-exercise muscle glycogen content. *Journal of Physiology* (in the Press).
- STEENBERG, A., TOFT, A. D., SCHJERLING, P., HALKJAER-KRISTENSEN, J. & PEDERSEN, B. K. (2001*b*). Plasma interleukin-6 during strenuous exercise: role of epinephrine. *American Journal of Physiology – Cell Physiology* **281**, C1001–1004.
- STEENBERG, A., VAN HALL, G., OSADA, T., SACCHETTI, M., SALTIN, B. & PEDERSEN, B. K. (2000). Production of interleukin-6 in contracting human skeletal muscles can account for the exercise-induced increase in plasma interleukin-6. *Journal of Physiology* **529**, 237–242.
- STEENBERG, A., VISSING, J. & PEDERSEN, B. K. (2001*c*). Lack of IL-6 production during exercise in patients with mitochondrial myopathy. *European Journal of Applied Physiology* **84**, 155–157.
- STOUTHARD, J. M., OUDE ELFERINK, R. P. & SAUERWEIN, H. P. (1996). Interleukin-6 enhances glucose transport in 3T3-L1 adipocytes. *Biochemical and Biophysical Research Communications* **220**, 241–245.
- STOUTHARD, J. M., ROMIJN, J. A., VAN DER POLL, T., ENDERT, E., KLAIN, S., BAKKER, P. J., VEENHOF, C. H. & SAUERWEIN, H. P. (1995). Endocrinologic and metabolic effects of interleukin-6 in humans. *American Journal of Physiology* **268**, E813–819.
- SUNDGREN-ANDERSSON, A. K., OSTLUND, P. & BARTFAI, T. (1998). IL-6 is essential in TNF- α -induced fever. *American Journal of Physiology* **275**, R2028–2034.
- TAKAI, Y., WONG, G. G., CLARK, S. C., BURAKOFF, S. J. & HERRMANN, S. H. (1988). B cell stimulatory factor-2 is involved in the differentiation of cytotoxic T lymphocytes. *Journal of Immunology* **140**, 508–512.
- TOFT, A. D., OSTROWSKI, K., ASP, S., MØLLER, K., IVERSEN, S., HERMANN, C., SØNDERGAARD, S. R. & PEDERSEN, B. K. (2000). The effects of n-3 PUFA on the cytokine response to strenuous exercise. *Journal of Applied Physiology* **89**, 2401–2405.
- TSIGOS, C., KYROU, I., CHALA, E., TSAPOGAS, P., STAVRIDIS, J. C., RAPTIS, S. A. & KATSIAMBROS, N. (1999). Circulating tumour necrosis factor α concentrations are higher in abdominal versus peripheral obesity. *Metabolism* **48**, 1332–1335.
- TSIGOS, C., PAPANICOLAOU, D. A., KYROU, I., DEFENSOR, R., MITSILADIS, C. S. & CHROUSOS, G. P. (1997). Dose-dependent effects of recombinant human interleukin-6 on glucose regulation. *Journal of Clinical Endocrinology and Metabolism* **82**, 4167–4170.
- TSUJINAKA, T., EBISUI, C., FUJITA, J., KISHIBUCHI, M., MORIMOTO, T., OGAWA, A., KATSUME, A., OHSUGI, Y., KOMINAMI, E. & MONDEN, M. (1995). Muscle undergoes atrophy in association with increase of lysosomal cathepsin activity in interleukin-6 transgenic mouse. *Biochemical and Biophysical Research Communications* **207**, 168–174.
- TSUJINAKA, T., FUJITA, J., EBISUI, C., YANO, M., KOMINAMI, E., SUZUKI, K., TANAKA, K., KATSUME, A., OHSUGI, Y., SHIOZAKI, H. & MONDEN, M. (1996). Interleukin 6 receptor antibody inhibits muscle atrophy and modulates proteolytic systems in interleukin 6 transgenic mice. *Journal of Clinical Investigation* **97**, 244–249.
- ULLUM, H., HAAHR, P. M., DIAMANT, M., PALMO, J., HALKJAER-KRISTENSEN, J. & PEDERSEN, B. K. (1994). Bicycle exercise enhances plasma IL-6 but does not change IL-1 α , IL-1 β , IL-6, or TNF- α pre-mRNA in BMNC. *Journal of Applied Physiology* **77**, 93–97.
- VAN WAGONER, N. J. & BENVENISTE, E. N. (1999). Interleukin-6 expression and regulation in astrocytes. *Journal of Neuroimmunology* **100**, 124–139.
- VGONTZAS, A. N., PAPANICOLAOU, D. A., BIXLER, E. O., HOPPER, K., LOTSIKAS, A., LIN, H. M., KALES, A. & CHROUSOS, G. P. (2000). Sleep apnea and daytime sleepiness and fatigue: relation to visceral obesity, insulin resistance, and hypercytokinemia. *Journal of Clinical Endocrinology and Metabolism* **85**, 1151–1158.
- VIERCK, J., O'REILLY, B., HOSSNER, K., ANTONIO, J., BYRNE, K., BUCCI, L. & DODSON, M. (2000). Satellite cell regulation following myotrauma caused by resistance exercise. *Cell Biology International* **24**, 263–272.
- YAN, S. F., TRITTO, I., PINSKY, D., LIAO, H., HUANG, J., FULLER, G., BRETT, J., MAY, L. & STERN, D. (1995). Induction of interleukin 6 (IL-6) by hypoxia in vascular cells. Central role of the binding site for nuclear factor-IL-6. *Journal of Biological Chemistry* **270**, 11463–11471.
- YUDKIN, J. S., KUMARI, M., HUMPHRIES, S. E. & MOHAMMED-ALI, V. (2000). Inflammation, obesity, stress and coronary heart disease: is interleukin-6 the link? *Atherosclerosis* **148**, 209–214.
- ZHANG, Y., PILON, G., MARETTE, A. & BARACOS, V. E. (2000). Cytokines and endotoxin induce cytokine receptors in skeletal muscle. *American Journal of Physiology – Endocrinology and Metabolism* **279**, E196–205.
- ZILBERSTEIN, A., RUGGIERI, R., KORN, J. H. & REVEL, M. (1986). Structure and expression of cDNA and genes for human interferon- β -2, a distinct species inducible by growth-stimulatory cytokines. *EMBO Journal* **5**, 2529–2537.

Acknowledgements

The original work from our group on IL-6 was supported by The Danish National Research Foundation (no. 504-14).

Corresponding author

B. K. Pedersen: Department of Infectious Diseases M7641, Rigshospitalet, Tagensvej 20, 2200 Copenhagen N, Denmark.

Email: bkp@rh.dk